

Ambulatory Care Pavilion- Level 2 Address: 3 Edmund D. Pellegrino Road, Stony Brook NY Our phone number is: 631-444-1759

Preparation for Esophageal Manometry with pH/Impedance ON MEDS

- 1. At least 24 Hours BEFORE the procedure:
- **<u>STOP</u>** the following medications if you are taking them for your stomach and esophageal problems.
 - Nitrates (Nitroglycerin, Nitro-patches/pastes, Isosorbide Dinitrate)
 - Calcium channel-blocking agents (Nifedipine, Verapamil, Diltiazem)
 - Anticholinergics (Propantheline)
 - Promotility agents (Metoclopramide, Bethanechol)
 - Muscle relaxants (Diazepam, Tizanidine, etc.)
 - Narcotics (Opiates)
 - Anti-Anxiety medications (Diazepam, Xanax, Ativan, etc.)
- Please continue to take any heart medication.

2. <u>CONTINUE</u> taking your acid medication (Prilosec, Nexium, Omeprazole, pantoprazole, Aciphex, Dexilant, Zantac, Pepcid, etc.).

SIX (6) HOURS BEFORE: <u>DO NOT</u> eat or drink anything, except for medication you are allowed to take. Take your acid medication with sips of water if they are due within this 6 hours.

4. DAY OF TEST:

- Stop at Patient Registration in the ACP building on Level 2 before proceeding to the "GI Motility Center".
- If you have a prescription order form, please bring it with you.
- We will be using medical adhesive above your lip/under your nose to help secure the Impedance probe to your face for 24 hours. We recommend for best results:
 - > Avoid use of cream and/or foundation/makeup on your face on the day of your appointment.
 - Shave off or trim beard.
 - > Bring or wear a zip up hoodie/jacket to wear over the Impedance box.

5. Please be prepared to spend up to 2 hours in the Motility Center.

6. You will need to return the following day to have the probe removed and return the equipment. This appointment will take 5 minutes. You do not need anyone to accompany you to either appointment.