

# Upper Endoscopy/ERCP Instructions

Name:	
Procedure:	
Physician:	
Location:	
Date of Procedure:	
Arrive at:	

**Please note:** As a courtesy to other patients and to your doctor, please call (631) 444-5220 as soon as possible if you are unable to keep your appointment.



## AS SOON AS POSSIBLE

1. Make arrangements for a responsible adult to accompany you when leaving the procedure, even if taking public transportation. Your procedure may be canceled if we cannot confirm your transportation on the day of the procedure. Taxi and Rideshare (Uber®, Lyft® etc.) drivers are not acceptable.

\*\*\*\***IMPORTANT**: IF YOU DO NOT HOLD CERTAIN MEDICATION(S) AS INSTRUCTED BELOW OR OBTAIN ANY NECESSARY CLEARANCES, YOUR PROCEDURE MAY BE CANCELED. IT IS YOUR RESPONSIBILITY THAT CLEARANCES ARE FAXED TO 631-444-5225 WITH DR.'S NAME AND DATE OF PROCEDURE

2. If you take any medications that increase your risk of bleeding, you must ask the prescribing doctor when to stop taking the medication before the procedure. The most common of these medications include but are not limited to:

Warfarin (Coumadin®)	Effient (Prasugrel®)	
Clopidogrel (Plavix®)	Apixaban (Eliquis®)	
Ticagrelor (Brilinta®)	Dabigatran (Pradaxa®)	
Rivaroxaban (Xarelto®)	Enoxaparin (Lovenox®)	

You may continue taking aspirin unless otherwise instructed

- 3. For patients with diabetes
  - If you take insulin, contact your diabetes doctor a minimum of **14 days prior to your procedure date** to adjust your insulin dose during the bowel preparation
  - If you use an insulin pump, a written letter from your endocrinologist giving clear instructions on adjustments for basal rate before and during the procedure will be required (the letter must be obtained at least 14 days prior to your procedure date)
  - If you use oral diabetes medications (except an SGLT-2 inhibitor), do not take these medications on the morning of your procedure; you may resume this medication once you resume a regular diet.
  - If you take an SGLT-2 inhibitor (Canigliflozin [Invokana], Dapagliflozin [Farxiga], or Empagliflozin [Jardiance]) for diabetes, do not take this medication for 3 days prior to your procedure date. If you take Ertugliflozin [Steglatro], do not take this medication for 4 days prior to your procedure date. After your procedure, you may take this medication once you resume a regular diet. If this medication is prescribed for a cardiac indication, please discuss holding this medication with your Cardiologist.



4. For patients taking a GLP-1 Agonist for diabetes or obesity

Drug	Interruption of Therapy	Resumption of Therapy
Once Daily GLP-1RA (SQ) Liraglutide (Victoza®, Saxenda®)	Hold day of procedure	Resume next day
Twice-Daily GLP-1RA (SQ) Exenatide (Byetta®)	Hold the night before procedure and the morning of the procedure	Resume next day/next dose
Once Daily GLP-1RA (Oral) Semaglutide (Rybelsus®)	Hold one week before the procedure	Resume next day
Weekly GLP-1 RA (SQ) Dulaglutide (Trulicity®) Exenatide (Bydureon ®) Semaglutide (Ozempic®, Wegovy®) Tirzapetide (Mounjaro®)	Hold one week before the procedure	Resume next day /next dose

5. If you have a pacemaker or defibrillator, please make sure your doctor is aware and bring the device card with you to your procedure



## **5 DAYS PRIOR TO YOUR UPPER ENDOSCOPY OR ERCP**

- Call the Admitting Office at (631) 444-1870 and provide your insurance information
- Confirm your transportation for the day of your procedure
- Please call if you need to cancel reschedule your procedure

### PREPARATION BEFORE YOUR UPPER ENDOSCOPY OR ERCP

- For safety reasons, you cannot have anything to eat or drink after midnight prior to the exam.
- If your exam is scheduled to start after 1PM, you may have small volumes of clear liquids until 7AM
- If you take medications for your heart, blood pressure or asthma, please take them with a sip of water at least 2 hours prior to the procedure. Do not take any oral diabetes medications.
- If you take a GLP-1 Receptor Agonist (see above section on medications), please **do not eat solids** for 12 hours before your procedure start time.

#### □ FOR PROCEDURES AT THE HOSPITAL

- Park in the main visitor parking garage; validation will be provided for one vehicle. Valet parking IS NOT eligible for validation. You must bring the parking ticket to endoscopy for validation.
- Report to registration on hospital level 5 (lobby) 1 hour before your scheduled procedure time. They will then send you to endoscopy on 14 North.

#### □ FOR PROCEDURES AT THE AMBULATORY CARE PAVILION (ACP)

- Directions and map to the ACP building can be found on the next page
- You can park at the gated parking lot in front of the building. You must bring the parking ticket to endoscopy for validation.
- Check in at the front desk 1 hour before your procedure and they will direct you to endoscopy

You should expect to be at the hospital or ACP for approximately 3 hours from time of registration to discharge from the endoscopy suite

#### The Ambulatory Care Pavilion at Stony Brook Medicine. 3 Edmund Pellegrino Road, Stony Brook NY

#### Directions:

- Heading north on Nicolls Road, turn right onto E Loop Road. Then, make a left at the first traffic light onto Edmund D Pellegrino Road.
- While on Edmund D Pellegrino Road, head towards the signs that say "Breast Center" (we are located in the same building as the Breast Center/Old Cancer Center).
- Drive through the 1<sup>st</sup> traffic circle and continue straight.
- At the 2<sup>nd</sup> traffic circle, make a U-turn so you are going the opposite direction (see attached map for details).
- Veer to the right and you will see an entrance to a small gated parking lot in front of the building.
- Please press the button and take a parking ticket that we will validate for you. If the gate is up, drive through it and park.
- The front desk will ask you for your name and will direct you to endoscopy

